arXiv:0809.3771v2 [math.AG] 29 Sep 2008

Test for reality of algebraic functions

S.M.Natanzon[1]

In this paper I prove the next affirmation

Theorem 1. *1) A complex algebraic function on a real algebraic curve with real points is equivalent to a real algebraic function, if and only if the divisor of preimage of its critical values is stable under the involution of complex conjugation. 2) A complex algebraic function on a real algebraic curve without real points is equivalent to a real or pseudoreal algebraic function, if and only if the divisor of preimage of its critical values is stable under the involution of complex conjugation.*

The proof uses representations of algebraic functions by Fuchsian groups as [N1, N(sect.1.6)]. It is interesting to compare this theorem with a conjecture of B. and M. Shapiro. It states, that a complex algebraic function is equivalent to real, if all its critical points are real. Proofs of special cases of this conjecture, when genus and degree satisfy $\deg(f)^2 - 4\deg(f) + 3 < 3g$ or $\deg(f) \leq 4$, are results of papers [EG1,EG2, ESS, DEISS].

Let us express our theorem in term of Riemann surfaces. A real algebraic curve is equivalent to a pair (P, τ), where P is a compact Riemann surface (the set of complex points of the curve) and $\tau : P \to P$ is an antiholomorphic involution (the involution of complex conjugation). Fixed points of τ are real points of the real algebraic curve[AG].

A complex algebraic function on a real curve (P, τ) is a holomorphic map $f : P \to \hat{\mathbb{C}} = \mathbb{C} \cup \infty$. The number ord_p is called the number of ramification of f in $p \in P$, if $f(z) = z^{\mathsf{ord}_p}$ in some local map $z : U \to \mathbb{C}$, where $z(p) = 0$. The points $\Sigma_P = \{p \in P | \mathsf{ord}_p > 1\}$ form the set of critical points. Its image $\Sigma_{\hat{\mathbb{C}}} = f(\Sigma_P) \subset \hat{\mathbb{C}}$ form the set of critical values. Let us consider the preimage of critical values $\Sigma_{P\hat{\mathbb{C}}} = f^{-1}(\Sigma_{\hat{\mathbb{C}}}) \subset P$ and its divisor $\Sigma(f) = \sum_{p\in\Sigma} \mathsf{ord}_p\, p$

Functions f and $\tilde{f}$ are called equivalent if $\tilde{f} = gf$ for $g \in \mathsf{Aut}(\hat{\mathbb{C}})$. A complex algebraic function is called real, if $\overline{f(\tau(p))} = f(p)$, and pseudoreal if $\overline{f(\tau(p))} = -\frac{1}{f(p)}$ [N2, N(sect.3.4)].

Let us illustrate this definitions by example. Consider a polynomial $F(x, y)$ with real coefficients. Then the affine real algebraic curve $F(x, y) = 0$ generates the pair (P_F, τ_F), where P_F is a regularization for complex points $\{(x, y) \in \mathbb{C}^2 | F(x, y) = 0\}$ of the curve and $\tau_F(x, y) = (\bar{x}, \bar{y})$. The set of real points of the real curve is the closure of $\{(x, y) \in \mathbb{R}^2 | F(x, y) = 0\}$. Complex algebraic functions are generated by restrictions of complex polynomials $G(x, y)$ on $\{(x, y) \in \mathbb{C}^2 | F(x, y) = 0\}$. The function is real if and only if the polynomial $G(x, y)$ has real coefficients.

Thus, theorem 1 is equivalent to

Theorem 2. *Let P be a compact Riemann surface, $f : P \to \hat{\mathbb{C}}$ be a holomorphic map, $\tau : P \to P$ be a antiholomorphic involution. Then $\tau\Sigma(f) = \Sigma(f)$, if an only if $\overline{gf\tau} = gf$ or $\overline{gf\tau} = -\frac{1}{gf}$ for $g \in \mathsf{Aut}(\hat{\mathbb{C}})$, and, moreover, $\overline{gf\tau} = gf$ for $\{p \in P | \tau p = p\} \neq \emptyset$.*

Proof. It is obviously that $\tau\Sigma(f) = \Sigma(f)$ follows from $\overline{gf\tau} = gf$ and $\overline{gf\tau} = -\frac{1}{gf}$. We use the induction by the degree $\deg(f)$ in order that to prove that $\tau\Sigma(f) = \Sigma(f)$ generate

---

[1] Supported by grants RFBR-04-01-00762, NWO 047.011.2004.026 (RFBR 05-02-89000-HBO-a), NSh-4719.2006.1, MTM 2005-01637, CRDF RM1-2543-MO-03, INTAS 05-7805

$\overline{gf\tau} = gf$ or $\overline{gf\tau} = -\frac{1}{gf}$. The theorem is obviously for $\deg(f) = 0$. Let $\deg(f) > 0$ and, thus, $\Sigma(f) \neq 0$. Consider an uniformisation $\Lambda \to S$ of $S = \hat{\mathbb{C}} \setminus f(\Sigma(f))$ by a Fuchsian group Γ on $\Lambda = \{z \in \mathbb{C} | \mathsf{Im(z)} > \mathsf{0}\}$. Then a subgroup $\tilde{\Gamma} \subset \Gamma$ uniformises the Riemann surface $\tilde{P} = P \setminus f^{-1}(f(\Sigma(f)))$. The involution τ is generated by an antiholomorphic map $\check{\tau} : \Lambda \to \Lambda$, where $\check{\tau}\tilde{\Gamma}\check{\tau}^{-1} = \tilde{\Gamma}$.

Consider $x \in \Sigma(f)$. Let $\check{x} \in \mathbb{R}$ be the fixed point of a parabolic automorphism $\tilde{\gamma} \in \tilde{\Gamma}$, that correspond to puncture x. Then the stationary subgroup $\tilde{\Gamma}_{\check{x}} = \{\gamma \in \tilde{\Gamma} | \gamma(\check{x}) = \check{x}\}$ is a subgroup of index $\mathsf{ord}_x(f) > 1$ of $\Gamma_{\check{x}} = \{\gamma \in \Gamma | \gamma(\check{x}) = \check{x}\}$. Thus, using $\tau\Sigma(f) = \Sigma(f)$ we have $\check{\tau}\tilde{\Gamma}_{\check{x}}\check{\tau}^{-1} \subset \tilde{\Gamma}$ and $\check{\tau}\Gamma_{\check{x}}\check{\tau}^{-1} \subset \Gamma$. Moreover, the condition $\tau\Sigma(f) = \Sigma(f)$ is equivalent to coincidence of indexes of stationer subgroups $[\Gamma_{\tilde{x}} : \tilde{\Gamma}_{\tilde{x}}] = [\Gamma_{\check{\tau}(\tilde{x})} : \tilde{\Gamma}_{\check{\tau}(\tilde{x})}]$ for all fixed points $\tilde{x}$ of parabolic automorphisms from $\tilde{\Gamma}$.

Let $\hat{\Gamma} \subset \Gamma$ be the subgroup, generated by $\tilde{\Gamma}$, $\Gamma_{\tilde{x}}$ and $\check{\tau}\Gamma_{\check{x}}\check{\tau}^{-1}$. It contains $\tilde{\Gamma}$ as a proper subgroup. Moreover, $\check{\tau}\hat{\Gamma}\check{\tau}^{-1} = \hat{\Gamma}$ and $[\Gamma_{\tilde{x}} : \hat{\Gamma}_{\tilde{x}}] = [\Gamma_{\hat{\tau}(\tilde{x})} : \hat{\Gamma}_{\hat{\tau}(\tilde{x})}]$ for all fixed points $\tilde{x}$ of parabolic automorphisms from $\hat{\Gamma}$.

Consider $\hat{P} = \Lambda/\hat{\Gamma}$. The inclusions of groups $\tilde{\Gamma} \subset \hat{\Gamma} \subset \Gamma$ generate the coverings $\Lambda \to \tilde{P} \xrightarrow{\tilde{\psi}} \hat{P} \xrightarrow{\hat{\psi}} S$. These coverings generate holomorphic maps of compact surfaces $P \xrightarrow{\varphi} \dot{P} \xrightarrow{\dot{\varphi}} \hat{\mathbb{C}}$ after gluing of punctures. Moreover, $f = \dot{\varphi}\varphi$, and $\deg(\dot{\varphi}) < \deg(f)$. The involution $\check{\tau}$ generates an antiholomorphic involution $\dot{\tau}$ on $\dot{P}$. Moreover, $\varphi\tau = \dot{\tau}\varphi$, and $\dot{\tau}\Sigma(\dot{\varphi}) = \Sigma(\dot{\varphi})$. Therefore, by inductive hypothesis $\overline{\dot{\varphi}\dot{\tau}}$ is equal to $g\dot{\varphi}$ or $-\frac{1}{g\dot{\varphi}}$ for $g \in \mathsf{Aut}(\hat{\mathbb{C}})$. Thus, the function $\overline{f\tau} = \overline{\dot{\varphi}\varphi\tau} = \overline{\dot{\varphi}\dot{\tau}\varphi}$ is equal $g\dot{\varphi}\varphi = gf$ or $-\frac{1}{g\dot{\varphi}\varphi} = -\frac{1}{gf}$. It is follow from [N2,N(sect.3.4)], that $\{p \in P | \tau p = p\} = \emptyset$ in the last case.

□

References

[AG] Alling N.L., Greenleaf N., Foundations of the theory of Klein surfaces. Berlin - Heidelbec rg - N.Y.: Springer - Verlag, 1971. Lecture Notes in Mathematics. V.219.

[EG1] Eremenko A., Gabrielov A., Elementary proof of the B. and M. Shapiro conjecture for rational functions, 2005, math/0512370v1, 21 p.

[EG2] Eremenko A., Gabrielov A., Rational functions with real critical points and the B. and M. Shapiro conjecture in real enumerative geometry. Ann. of Math. (2) 155 (2002), no.1, 105-129.

[ESS] Ekedahl T, Shapiro B., Shapiro M., First steps towards total reality of meromorphic functions, Mosc. Math. J. 6 (2006), no.1, 95-106.

[DEISS] Degtyarev A., Ekedahl T., Itenberg I., Shapiro B., Shapiro M., On total reality of meromorphic functions, Ann. Inst. Fourier (Grenoble) 57 (2007), no. 6, 2015-2030 (math/0605077).

[N1] Natanzon S.M., Uniformization of spaces of meromorphic functions. Dokl. Acad. Nauk SSSR, 287:5 (1986), 1058-1061 (Russian); English: Soviet. Math. Dokl., 33:2(1986), 487-490.

[N2] Natanzon S.M., Topology of 2-dimensional coverings and meromorphic functions on real and complex algebraic curves. Selecta Math.Soviet., 12:3 (1993), 251-291.

[N] Natanzon S.M., Moduli of Riemann surfaces, real algebraic curves, and their superanalogs. Translations of Mathematical Monographs, American Mathematical Society, Vol.225, (2004), 160 P.